Quasi-bound states of massive scalar fields in the Kerr black-hole spacetime: Beyond the hydrogenic approximation

Shahar Hod
The Ruppin Academic Center, Emeq Hefer 40250, Israel
and
The Hadassah Institute, Jerusalem 91010, Israel
(Dated: October 21, 2015)

Rotating black holes can support quasi-stationary (unstable) bound-state resonances of massive scalar fields in their exterior regions. These spatially regular scalar configurations are characterized by instability timescales which are much longer than the timescale M set by the geometric size (mass) of the central black hole. It is well-known that, in the small-mass limit $\alpha \equiv M\mu \ll 1$ (here μ is the mass of the scalar field), these quasi-stationary scalar resonances are characterized by the familiar *hydrogenic* oscillation spectrum: $\omega_{\rm R}/\mu = 1-\alpha^2/2\bar{n}_0^2$, where the integer $\bar{n}_0(l,n;\alpha\to 0) = l+n+1$ is the principal quantum number of the bound-state resonance (here the integers $l = 1,2,3,...$ and $n = 0,1,2,...$ are the spheroidal harmonic index and the resonance parameter of the field mode, respectively). As it depends only on the principal resonance parameter $\bar{n}_0$, this *small*-mass ($\alpha \ll 1$) hydrogenic spectrum is obviously *degenerate*. In this paper we go beyond the small-mass approximation and analyze the quasi-stationary bound-state resonances of massive scalar fields in rapidly-spinning Kerr black-hole spacetimes in the regime $\alpha = O(1)$. In particular, we derive the non-hydrogenic (and, in general, *non*-degenerate) resonance oscillation spectrum $\omega_{\rm R}/\mu = \sqrt{1-(\alpha/\bar{n})^2}$, where $\bar{n}(l,n;\alpha) = \sqrt{(l+1/2)^2-2m\alpha+2\alpha^2}+1/2+n$ is the *generalized* principal quantum number of the quasi-stationary resonances. This analytically derived formula for the characteristic oscillation frequencies of the composed black-hole-massive-scalar-field system is shown to agree with direct numerical computations of the quasi-stationary bound-state resonances.

arXiv:1510.05649v1 [gr-qc] 19 Oct 2015

I. INTRODUCTION

Recent analytical [1] and numerical [2] studies of the coupled Einstein-scalar equations have revealed that rotating black holes can support stationary spatially regular configurations of *massive* scalar fields in their exterior regions. These bound-state resonances of the composed black-hole-scalar-field system owe their existence to the well-known phenomenon of superradiant scattering [3, 4] of integer-spin (bosonic) fields in rotating black-hole spacetimes.

The stationary black-hole-scalar-field configurations [1, 2] mark the boundary between stable and unstable bound-state resonances of the composed system. In particular, these stationary scalar field configurations are characterized by azimuthal frequencies $\omega_{\rm field}$ which are in resonance with the black-hole angular velocity $\Omega_{\rm H}$ [5]:

$$\omega_{\rm field} = m\Omega_{\rm H}\ , \tag{1}$$

where $m = 1,2,3,...$ is the azimuthal quantum number of the field mode. Bound-state field configurations in the superradiant regime $\omega_{\rm field} < m\Omega_{\rm H}$ are known to be unstable (that is, grow in time), whereas bound-state field configurations in the regime $\omega_{\rm field} > m\Omega_{\rm H}$ are known to be stable (that is, decay in time) [3, 4].

The bound-state scalar resonances of the rotating Kerr black-hole spacetime are characterized by at least two different time scales: (1) the typical oscillation period $\tau_{\rm oscillation} \equiv 2\pi/\omega_{\rm R} \sim 1/\mu$ of the bound-state massive scalar configuration (here μ is the mass of the scalar field [6]), and (2) the instability growth time scale $\tau_{\rm instability} \equiv 1/\omega_{\rm I}$ associated with the superradiance phenomenon. Former studies [7–11] of the Einstein-massive-scalar-field system have revealed that these two time scales are well separated. In particular, it was shown [7–11] that the composed system is characterized by the relation

$$\tau_{\rm instability} \gg \tau_{\rm oscillation}\ , \tag{2}$$

or equivalently

$$\omega_{\rm I} \ll \omega_{\rm R}\ . \tag{3}$$

The strong inequality (2) implies that the bound-state massive scalar configurations may be regarded as the *quasi-stationary* resonances of the composed system.

As shown in [12, 13], the physical significance of the characteristic black-hole-scalar-field oscillation frequencies $\{\omega_{\rm R}(n)\}_{n=0}^{n=\infty}$ [14] lies in the fact that the corresponding quasi-stationary scalar resonances dominate the dynamics of massive scalar fields in curved black-hole spacetimes. In particular, recent numerical simulations [12, 13] of the

dynamics of massive scalar fields in the Kerr black-hole spacetime have demonstrated explicitly that these quasi-stationary bound-state resonances dominate the characteristic Fourier power spectra $P(\omega)$ of the composed black-hole-massive-scalar-field system [15].

II. THE SMALL-MASS HYDROGENIC SPECTRUM

As shown by Detweiler [7], the massive scalar resonances can be calculated analytically in the small-mass regime $M\mu \ll 1$. In particular, one finds [7] that the quasi-stationary bound-state scalar resonances are characterized by the familiar *hydrogenic* spectrum

$$\frac{\omega_{\rm R}(\bar{n}_0)}{\mu}=1-\frac{\alpha^2}{2\bar{n}_0^2}\ \ \ \ \text{for}\ \ \ \ \alpha\equiv M\mu\ll 1\ , \tag{4}$$

where the integer

$$\bar{n}_0(l,n;\alpha\to 0)=l+1+n \tag{5}$$

is the principal quantum number of the quasi bound-state resonances. Here the integer $l\geq |m|$ is the spheroidal harmonic index of the field mode and $n=0,1,2,...$ is the resonance parameter.

It is worth emphasizing that the hydrogenic spectrum (4) depends only on the principal resonance parameter (quantum number) $\bar{n}_0=l+1+n$. This small-mass oscillation spectrum is therefore *degenerate*. That is, two different modes, (l,n) and (l',n') with $l+n=l'+n'$, are characterized by the *same* resonant frequency: $\omega_{\rm R}(l,n)=\omega_{\rm R}(l',n')$ for $l+n=l'+n'$.

To the best of our knowledge, the oscillation frequency spectrum which characterizes the quasi-stationary bound-state resonances of massive scalar fields in the rotating Kerr black-hole spacetime has not been studied analytically beyond the hydrogenic regime (4) of small ($\alpha\ll 1$) field masses. The main goal of the present paper is to analyze the oscillation spectrum of the composed black-hole-massive-scalar-field system in the $\alpha=O(1)$ regime. To that end, we shall use the resonance equation [see Eq. (11) below] derived in [10] for the bound-state resonances of massive scalar fields in rapidly-rotating (near-extremal) Kerr black-hole spacetimes. As we shall show below, this resonance equation can be solved *analytically* to yield the characteristic oscillation spectrum $\{\omega_{\rm R}(n)\}_{n=0}^{n=\infty}$ of the quasi-stationary bound-state scalar resonances in the regime $\alpha\lesssim 1$.

III. DESCRIPTION OF THE SYSTEM

We consider a scalar field Ψ of mass μ linearly coupled to a rapidly-rotating (near-extremal) Kerr black hole of mass M and dimensionless angular momentum $a/M\to 1^-$. The dynamics of the scalar field in the black-hole spacetime is governed by the Klein-Gordon (Teukolsky) wave equation

$$(\nabla^a\nabla_a-\mu^2)\Psi=0\ . \tag{6}$$

Substituting the field decomposition [16–18]

$$\Psi(t,r,\theta,\phi)=\int\sum_{l,m}e^{im\phi}S_{lm}(\theta)R_{lm}(r)e^{-i\omega t}d\omega \tag{7}$$

into the wave equation (6), one finds [19, 20] that the radial function R and the angular function S obey two ordinary differential equations of the confluent Heun type [21, 22].

The angular eigenfunctions, known as the spheroidal harmonics, are determined by the angular Teukolsky equation [19–23]

$$\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\Big(\sin\theta\frac{\partial S_{lm}}{\partial\theta}\Big)+\Big[a^2(\omega^2-\mu^2)\cos^2\theta-\frac{m^2}{\sin^2\theta}+A_{lm}\Big]S_{lm}=0\ . \tag{8}$$

The regularity requirements of these functions at the two boundaries $\theta=0$ and $\theta=\pi$ single out a discrete set of angular eigenvalues $\{A_{lm}\}$ [see Eq. (14) below] labeled by the integers l and m.

The radial Teukolsky equation is given by [19, 20, 24]

$$\Delta\frac{d}{dr}\Big(\Delta\frac{dR_{lm}}{dr}\Big)+\Big\{[(r^2+a^2)\omega-am]^2-\Delta(a^2\omega^2-2ma\omega+\mu^2r^2+A_{lm})\Big\}R_{lm}=0\ , \tag{9}$$

where $\Delta \equiv (r-r_+)(r-r_-)$ [25]. Note that the angular Teukolsky equation (8) and the radial Teukolsky equation (9) are coupled by the angular eigenvalues $\{A_{lm}\}$.

The quasi-stationary bound-state resonances of the massive scalar fields in the black-hole spacetime are characterized by the boundary conditions of purely ingoing waves at the black-hole horizon (as measured by a comoving observer) and a spatially decaying (bounded) radial eigenfunction at asymptotic infinity [7–9, 26]:

$$R_{lm} \sim \begin{cases} \frac{1}{r}e^{-\sqrt{\mu^2-\omega^2}\,y} & \text{as } r\to\infty \ \ (y\to\infty)\ ; \\ e^{-i(\omega-m\Omega_{\rm H})y} & \text{as } r\to r_+ \ \ (y\to-\infty)\ , \end{cases} \tag{10}$$

where $\Omega_{\rm H}$ is the angular velocity of the black-hole horizon [see Eq. (12) below]. The boundary conditions (10) imposed on the radial eigenfunctions single out a discrete set of eigenfrequencies $\{\omega_n(a/M,l,m,\alpha)\}_{n=0}^{n=\infty}$ which characterize the quasi-stationary bound-state resonances of the massive scalar fields in the Kerr black-hole spacetime [7–9].

IV. THE CHARACTERISTIC RESONANCE EQUATION AND ITS REGIME OF VALIDITY

Solving analytically the radial Klein-Gordon (Teukolsky) equation (9) in two different asymptotic regions and using a standard matching procedure for these two radial solutions in their common overlap region [see Eq. (15) below], we have derived in [10] the characteristic resonance equation

$$\frac{1}{\Gamma(\frac{1}{2}+\beta-\kappa)} = \Big[\frac{\Gamma(-2\beta)}{\Gamma(2\beta)}\Big]^2 \frac{\Gamma(\frac{1}{2}+\beta-ik)}{\Gamma(\frac{1}{2}-\beta-ik)\Gamma(\frac{1}{2}-\beta-\kappa)}\Big[8iMr_+\sqrt{\mu^2-\omega^2}(m\Omega_{\rm H}-\omega)\Big]^{2\beta} \tag{11}$$

for the bound-state resonances of the composed Kerr-black-hole-massive-scalar-field system. Here

$$\Omega_{\rm H} = \frac{a}{r_+^2+a^2} \tag{12}$$

is the angular velocity of the black-hole horizon, and

$$k \equiv 2\omega r_+ \quad , \quad \kappa \equiv \frac{\omega k-\mu^2 r_+}{\sqrt{\mu^2-\omega^2}} \quad , \quad \beta^2 \equiv a^2\omega^2 - 2ma\omega + \mu^2 r_+^2 + A_{lm} - k^2 + \frac{1}{4}\ , \tag{13}$$

where $\{A_{lm}\}$ are the angular eigenvalues which couple the radial Teukolsky equation (9) to the angular (spheroidal) equation (8). These angular eigenvalues can be expanded in the form [23]

$$A_{lm} = l(l+1) + \sum_{k=1}^{\infty} c_k a^{2k}(\mu^2-\omega^2)^k\ , \tag{14}$$

where the expansion coefficients $\{c_k\}$ are given in [23].

Before proceeding, it should be emphasized that the validity of the resonance equation (11) is restricted to the regime

$$\tau \ll M(m\Omega_{\rm H}-\omega) \ll x_{\rm o} \ll \frac{1}{M\sqrt{\mu^2-\omega^2}}\ , \tag{15}$$

where $\tau \equiv (r_+-r_-)/r_+ \ll 1$ is the dimensionless temperature of the rapidly-rotating (near-extremal) Kerr black hole, and the dimensionless coordinate $x_{\rm o} \equiv (r_{\rm o}-r_+)/r_+$ belongs to the *overlap* region in which the two different solutions of the radial Teukolsky equation (hypergeometric and confluent hypergeometric radial wave functions) can be matched together, see [10, 27] for details. The inequalities in (15) imply that the resonance condition (11) should be valid in the regime [28]

$$M^2(m\Omega_{\rm H}-\omega)\sqrt{\mu^2-\omega^2} \ll 1\ . \tag{16}$$

V. THE QUASI-STATIONARY BOUND-STATE RESONANCES OF THE COMPOSED BLACK-HOLE-MASSIVE-SCALAR-FIELD SYSTEM

As we shall now show, the resonance condition (11) can be solved *analytically* in the physical regime (16). In particular, in the present section we shall derive a (remarkably simple) analytical formula for the discrete spectrum

of oscillation frequencies, $\{\omega_{\rm R}(l,m,\alpha;n)\}_{n=0}^{n=\infty}$, which characterize the quasi-stationary bound-state resonances of the composed Kerr-black-hole-massive-scalar-field system.

Our analytical approach is based on the fact that the right-hand-side of the resonance equation (11) is small in the regime (16) with $\beta\in\mathbb{R}$ [29]. The resonance condition can therefore be approximated by the simple zeroth-order equation

$$\frac{1}{\Gamma(\frac{1}{2}+\beta-\kappa)}=0\ . \tag{17}$$

As we shall now show, this zeroth-order resonance condition can be solved analytically to yield the real oscillation frequencies which characterize the bound-state scalar resonances. We first use the well-known pole structure of the Gamma functions [23] in order to write the resonance equation (17) in the form [10]

$$\frac{1}{2}+\beta-\kappa=-n, \tag{18}$$

where the integer $n=0,1,2,...$ is the resonance parameter of the field mode.

Defining the dimensionless variable [30]

$$\epsilon\equiv M\sqrt{\mu^2-\omega^2}\ , \tag{19}$$

one finds from Eq. (13)

$$\beta^2=\beta_0^2+O(\epsilon^2,\tau)\quad\text{and}\quad\kappa=\frac{\alpha^2}{\epsilon}-2\epsilon+O(\tau)\ , \tag{20}$$

where

$$\beta_0^2\equiv(l+1/2)^2-2m\alpha-2\alpha^2\ . \tag{21}$$

Substituting (20) into the resonance condition $\beta^2=[\kappa-(n+1/2)]^2$ [see Eq. (18)], one obtains the characteristic equation

$$\epsilon^2\cdot[(2l+1)^2-8m\alpha+8\alpha^2-(2n+1)^2]+\epsilon\cdot4(2n+1)\alpha^2-4\alpha^4+O(\tau,\epsilon^3)=0 \tag{22}$$

for the dimensionless parameter ϵ. This resonance equation can easily be solved to yield

$$\epsilon(l,m;n)=\frac{2\alpha^2}{\sqrt{(2l+1)^2-8m\alpha+8\alpha^2}+1+2n}\ . \tag{23}$$

Finally, taking cognizance of the relation (19), one finds the discrete spectrum of oscillation frequencies

$$\frac{\omega_{\rm R}(n)}{\mu}=\sqrt{1-\Big(\frac{\alpha}{\ell+1+n}\Big)^2} \tag{24}$$

which characterize the quasi-stationary bound-state resonances of the composed black-hole-massive-scalar-field system. Here

$$\ell\equiv\frac{1}{2}\big[\sqrt{(2l+1)^2-8m\alpha+8\alpha^2}-1\big] \tag{25}$$

is the generalized (finite-mass) spheroidal harmonic index. Note that $\ell\to l$ in the small mass $\alpha\ll1$ limit, in which case one recovers from (24) the well-known hydrogenic spectrum (4) of [7].

It is worth noting that, in general, the parameter $\ell(\alpha)$ is not an integer. This implies that, for generic values of the dimensionless mass parameter α, the non-hydrogenic oscillation spectrum (24) is *not* degenerate [31].

VI. NUMERICAL CONFIRMATION

It is of physical interest to test the accuracy of the analytically derived formula (24) for the characteristic oscillation frequencies $\omega_{\rm R}^{\rm ana}(n)/\mu$ of the quasi-stationary massive scalar configurations. The quasi bound-state resonances can be computed using standard numerical techniques, see [9, 12] for details. In Table I we present a comparison between the

Resonance parameter n	0	1	2	3	4
$\omega_{\rm R}^{\rm ana}(n)/\omega_{\rm R}^{\rm num}(n)$	0.9999	1.0006	1.0004	1.0002	0.9994

TABLE I: Quasi-stationary resonances of massive scalar fields in the rotating Kerr black-hole spacetime. The data shown is for the fundamental $l=m=1$ mode with $a/M=0.99$, $\alpha\equiv M\mu=0.42$, and $n=0,1,2,3,4$. We display the dimensionless ratio between the *analytically* derived oscillation frequencies $\omega_{\rm R}^{\rm ana}(n)$ [see Eq. (24)] and the *numerically* computed resonances $\omega_{\rm R}^{\rm num}(n)$ of [12]. One finds a good agreement between the analytical formula (24) and the numerical data of [12] .

analytically derived oscillation frequencies (24) and the numerically computed resonances [12]. The data presented is for the fundamental $l=m=1$ mode with $a/M=0.99$ [32] and $\alpha=0.42$ [33, 34]. One finds a good agreement between the *analytically* calculated oscillation frequencies (24) and the *numerically* computed resonances of [12].

In order to compare the accuracy of the newly derived analytical formula (24) with the accuracy of the familiar hydrogenic (small-mass) spectrum (4), we display in Table II the physical quantity $\epsilon(n)$ [see Eq. (19)] which provides a quantitative measure for the deviation of the resonant oscillation frequency $\omega_{\rm R}(n)$ from the field mass parameter μ. In particular, we present the dimensionless ratios $\epsilon^{\rm ana}/\epsilon^{\rm num}$ and $\epsilon^{\text{ana-hydro}}/\epsilon^{\rm num}$, where $\epsilon^{\rm ana}(n)$ is given by the analytical formula (24), $\epsilon^{\text{ana-hydro}}(n)$ is defined from the hydrogenic spectrum (4), and $\epsilon^{\rm num}(n)$ is obtained from the numerically computed resonances of [12]. One finds that, in general, the newly derived formula (24) performs better than the hydrogenic formula (4) [35].

Resonance parameter n	0	1	2	3	4
$\epsilon^{\rm ana}(n)/\epsilon^{\rm num}(n)$	1.001	0.974	0.971	0.974	1.140
$\epsilon^{\text{ana-hydro}}(n)/\epsilon^{\rm num}(n)$	0.910	0.916	0.928	0.939	1.107

TABLE II: Quasi-stationary resonances of massive scalar fields in the rotating Kerr black-hole spacetime. The data shown is for the fundamental $l=m=1$ mode with $a/M=0.99$, $\alpha\equiv M\mu=0.42$, and $n=0,1,2,3,4$. We display the dimensionless ratios $\epsilon^{\rm ana}/\epsilon^{\rm num}$ and $\epsilon^{\text{ana-hydro}}/\epsilon^{\rm num}$, where $\epsilon^{\rm ana}(n)$ is given by the analytical formula (24), $\epsilon^{\text{ana-hydro}}(n)$ is defined from the hydrogenic spectrum (4), and $\epsilon^{\rm num}(n)$ is obtained from the numerically computed resonances of [12]. One finds that, in general, the newly derived analytical formula (24) performs better than the hydrogenic formula (4) [35].

VII. SUMMARY

In summary, we have studied the resonance spectrum of quasi-stationary massive scalar configurations linearly coupled to a near-extremal (rapidly-rotating) Kerr black-hole spacetime. In particular, we have derived a compact analytical expression [see Eq. (24)] for the characteristic oscillation frequencies $\omega_{\rm R}^{\rm ana}(n)/\mu$ of the bound-state massive scalar fields. It was shown that the *analytically* derived formula (24) agrees with direct *numerical* computations [12] of the black-hole-scalar-field resonances.

It is well known that the characteristic hydrogenic spectrum (4) in the *small*-mass $\alpha\ll 1$ limit is highly degenerate – it depends only on the principal resonance parameter $\bar{n}_0\equiv l+1+n$ [36] [Thus, according to (4), two different modes which are characterized by the integer parameters (l,n) and (l',n') with $l+n=l'+n'$ share the *same* resonant frequency $\omega_{\rm R}(\bar{n}_0)$ in the $\alpha\to 0$ limit]. On the other hand, the newly derived resonance spectrum (24), which is valid in the $\alpha=O(1)$ regime, is no longer degenerate. That is, for generic values of the dimensionless mass parameter α, two quasi-stationary modes with different sets of the integer parameters (l,n) are characterized, according to (24), by *different* oscillation frequencies [37].

Finally, it is worth emphasizing again that the physical significance of the characteristic oscillation frequencies (24) lies in the fact that these quasi-stationary (long-lived) resonances dominate the dynamics [and, in particular, dominate the characteristic Fourier power spectra $P(\omega)$ [12]] of the massive scalar fields in the black-hole spacetime.

ACKNOWLEDGMENTS

This research is supported by the Carmel Science Foundation. I would like to thank Yael Oren, Arbel M. Ongo,

Ayelet B. Lata, and Alona B. Tea for helpful discussions.

---

[1] S. Hod, Phys. Rev. D **86**, 104026 (2012) [arXiv:1211.3202]; S. Hod, The Euro. Phys. Journal C **73**, 2378 (2013) [arXiv:1311.5298]; S. Hod, Phys. Rev. D **90**, 024051 (2014) [arXiv:1406.1179]; S. Hod, Phys. Lett. B **739**, 196 (2014) [arXiv:1411.2609]; S. Hod, Class. and Quant. Grav. **32**, 134002 (2015); S. Hod, Phys. Lett. B **708**, 320 (2012) [arXiv:1205.1872].

[2] C. A. R. Herdeiro and E. Radu, Phys. Rev. Lett. **112**, 221101 (2014); C. A. R. Herdeiro and E. Radu, Phys. Rev. D **89**, 124018 (2014); C. A. R. Herdeiro and E. Radu, Int. J. Mod. Phys. D **23**, 1442014 (2014); C. L. Benone, L. C. B. Crispino, C. Herdeiro, and E. Radu, Phys. Rev. D **90**, 104024 (2014); C. Herdeiro, E. Radu, and H. Runarsson, Phys. Lett. B **739**, 302 (2014); C. Herdeiro and E. Radu, arXiv:1501.04319; C. A. R. Herdeiro and E. Radu, arXiv:1504.08209; C. A. R. Herdeiro and E. Radu, arXiv:1505.04189; J. C. Degollado and C. A. R. Herdeiro, Gen. Rel. Grav. **45**, 2483 (2013).

[3] Ya. B. Zel'dovich, Pis'ma Zh. Eksp. Teor. Fiz. **14**, 270 (1971) [JETP Lett. **14**, 180 (1971)]; Zh. Eksp. Teor. Fiz. **62**, 2076 (1972) [Sov. Phys. JETP **35**, 1085 (1972)]; A. V. Vilenkin, Phys. Lett. B **78**, 301 (1978).

[4] W. H. Press and S. A. Teukolsky, Nature **238**, 211 (1972); W. H. Press and S. A. Teukolsky, Astrophys. J. **185**, 649 (1973).

[5] We use natural units in which $G = c = \hbar = 1$.

[6] Note that the field mass parameter μ stands for $\mu/\hbar$. Hence, it has the dimensions of $(\text{length})^{-1}$.

[7] S. Detweiler, Phys. Rev. D **22**, 2323 (1980).

[8] T. M. Zouros and D. M. Eardley, Annals of physics **118**, 139 (1979).

[9] S. R. Dolan, Phys. Rev. D **76**, 084001 (2007).

[10] S. Hod and O. Hod, Phys. Rev. D **81**, 061502 Rapid communication (2010) [arXiv:0910.0734].

[11] R. Brito, V. Cardoso, and P. Pani, Class. Quant. Grav. **32**, 134001 (2015).

[12] S. Dolan, Phys. Rev. D **87**, 124026 (2013).

[13] H. Witek, V. Cardoso, A. Ishibashi, and U. Sperhake, Phys. Rev. D **87**, 043513 (2013).

[14] Here the integer $n = 0, 1, 2, ...$ is the resonance parameter of the radial field mode, see Eq. (18) below.

[15] More explicitly, by recording the temporal dependence of the scalar-field amplitude $A(t)$ at some fixed point $r = r_0$, and taking the Fourier transform of this time-dependent field amplitude, one finds [12] that the resulting Fourier power spectrum $P(\omega)$ is characterized by sharp picks at the appropriate resonant frequencies $\omega_{\rm R}(\mu)$ [see Eq. (24) below] of the composed black-hole-scalar-field system (see, in particular, Fig. 12 of [12]). This implies that the quasi-stationary bound-state scalar resonances are excited by generic initial data of the massive scalar fields.

[16] Here (t, r, θ, ϕ) are the Boyer-Lindquist coordinates [17].

[17] R. P. Kerr, Phys. Rev. Lett. **11**, 237 (1963); R. H. Boyer and R. W. Lindquist, J. Math. Phys. **8**, 265 (1967).

[18] Here ω, l, and m are respectively the (conserved) frequency of the field mode, its spheroidal harmonic index, and its azimuthal harmonic index.

[19] S. A. Teukolsky, Phys. Rev. Lett. **29**, 1114 (1972); S. A. Teukolsky, Astrophys. J. **185**, 635 (1973).

[20] T. Hartman, W. Song, and A. Strominger, JHEP 1003:118 (2010).

[21] A. Ronveaux, *Heun's differential equations*. (Oxford University Press, Oxford, UK, 1995); C. Flammer, *Spheroidal Wave Functions* (Stanford University Press, Stanford, 1957).

[22] P. P. Fiziev, e-print arXiv:0902.1277; R. S. Borissov and P. P. Fiziev, e-print arXiv:0903.3617; P. P. Fiziev, Phys. Rev. D **80**, 124001 (2009); P. P. Fiziev, Class. Quant. Grav. **27**, 135001 (2010).

[23] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions* (Dover Publications, New York, 1970).

[24] S. Hod, Phys. Rev. Lett. **100**, 121101 (2008) [arXiv:0805.3873]; S. Hod, Phys. Rev. D **75**, 064013 (2007) [arXiv:gr-qc/0611004].

[25] Here $r_\pm = M \pm (M^2 - a^2)^{1/2}$ are the black-hole (event and inner) horizons.

[26] Here the "tortoise" radial coordinate y is defined by the relation $dy = [(r^2 + a^2)/\Delta]dr$.

[27] J. G. Rosa, JHEP 1006:015 (2010).

[28] In particular, since each inequality in (15) roughly corresponds to an order-of-magnitude difference between two quantities, one concludes that the resonance condition (11) should be valid in the regime $M^2(m\Omega_{\rm H} - \omega)\sqrt{\mu^2 - \omega^2} \lesssim 10^{-2}$.

[29] One finds that $\beta \in \mathbb{R}$ in the regime $\alpha \leq [\sqrt{2(l + 1/2)^2 + m^2} - m]/2$, see Eq. (21) below.

[30] This physical parameter provides a quantitative measure for the deviation of the resonant oscillation frequency $\omega_{\rm R}(n)$ from the field mass parameter μ.

[31] Compare this property of the non-hydrogenic spectrum (24) with the familiar hydrogenic spectrum (4) which is known to be *degenerate*.

[32] This is the largest black-hole spin for which we have exact (numerical) data [12] for the characteristic oscillation frequencies $\omega_{\rm R}(n)/\mu$ of the quasi-stationary bound-state resonances. Note that the numerical results of [9] indicate that, for $\alpha \lesssim 0.6$ and $a \to 1^-$, the values of the fundamental resonant frequencies $\omega_{\rm R}/\mu$ depend very weakly on the black-hole rotation parameter a (see, in particular, Fig. 5 of [9]).

[33] As discussed above, our resonance equation (11) is restricted to the regime $M^2(m\Omega_{\rm H} - \omega)\sqrt{\mu^2 - \omega^2} \lesssim 10^{-2}$ [28]. Substituting our solution (24) into the left-hand-side of (16), one finds that, for the fundamental $l = m = 1$ mode, the requirement $M^2(m\Omega_{\rm H} - \omega)\sqrt{\mu^2 - \omega^2} \lesssim 10^{-2}$ is respected in the regime $\alpha \lesssim 0.58$. Note that this mass regime is consistent with our previous requirement $\beta \in \mathbb{R}$ [29], which implies the weaker restriction $\alpha \lesssim 0.67$ for the fundamental $l = m = 1$ mode.

[34] Note that Eqs. (21) and (23) yield $\beta_0^2 \simeq 1.057$ and $\epsilon(n=0) \simeq 0.097$ for the fundamental $l=m=1$ mode with $\alpha=0.42$. One therefore finds the characteristic small ratio $\epsilon^2/\beta_0^2 \simeq 0.009 \ll 1$. This strong inequality justifies our previous assumption $\epsilon^2/\beta_0^2 \ll 1$ [see Eq. (20)]. Moreover, since $\epsilon(n)$ is a *decreasing* function of the resonance parameter n [see Eq. (23)], one finds that the approximation $\epsilon^2/\beta_0^2 \ll 1$ becomes even better for the excited ($n \geq 1$) resonant modes.
[35] The only exception is the excited $n=4$ mode, for which the agreement between the numerically computed deviation-parameter $\epsilon^{\rm num}(n=4)$ and the analytical formulas (4) and (24) is quite poor.
[36] Note that this parameter is an integer.
[37] Note that, for generic values of the mass parameter α, the generalized (finite-mass) spheroidal harmonic index $\ell(\alpha)$ that appears in the resonance spectrum (24) is not an integer.